PATIENT INFORMATION: ZACCUR ISAAC

REPORT STATUS: FINAL

SPECIMEN INFORMATION ORDERING PHYSICIAN: WIN NGUYEN

DOB: 02/14/1992

SPECIMEN: BLOOD

AGE: 32

CLIENT INFORMATION

GENDER: MALE

LAB REF NO: SLV245789 COLLECTED: 10/18/2024

RECEIVED: 10/21/2024



REPORTED: 10/24/2024

#### Test Name Result

#### HEPATITIS B SURFACE ANTIGEN W/REFL CONFIRM

HEPATITIS B SURFACE ANTIGEN NON-REACTIVE

CONFIRMATION DNR

HEPATITIS C ANTIBODY

HEPATITIS C ANTIBODY NON-REACTIVE

SIGNAL TO CUT-OFF 0.01

#### HSV 1/2 IGG, TYPE SPECIFIC AB HERPESELECT

HSV 1 IGG TYPE SPECIFIC AB 0.01
HSV 2 IGG TYPE SPECIFIC AB 0.04

Value Interpretation
----<0.90 Negative

0.90-1.10 Equivocal >1.10 Positive

This assay utilizes recombinant type-specific antigens to differentiate HSV-1 from HSV-2 infections. A positive result cannot distinguish between recent and past infection. If recent HSV infection is suspected but the results are negative or equivocal, the assay should be repeated in 4-6 weeks. The performance characteristics of the assay have not been established for pediatric populations, immunocompromised patients, or neonatal screening.

## HIV AB, HIV 1/2, EIA, WITH REFLEXES

HIV 1/2 EIA AB SCREEN

NON-REACTIVE

A Nonreactive HIV-1/2 antibody result does not exclude HIV infection since the time frame for seroconversion is variable. If acute HIV infection is suspected, antibody retesting and nucleic acid amplification (HIV DNA/RNA) testing is recommended.

This assay exhibits increased sensitivity over the HIV-1 Western Blot test and simultaneously differentiates HIV-1 antibody reactivity from HIV-2 antibody reactivity.

## CHLAMYDIA/N. GONORRHOEAE RNA, TMA

CHLAMYDIA TRACHOMATIS RNA, TMA NOT DETECTED
NEISSERIA GONORRHOEAE RNA, TMA NOT DETECTED

COMMENT

RPR (DX) W/REFL TITER AND NON-REACTIVE

NON-REACTIVE ΤP

# Performing Laboratory Information:

Total Men's Primary Care- Preston Plano 1921 Preston Rd Suite B2076, Plano, TX 75093 (469) 750-1682

2 of 2